



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Ripley. The first, consisting of irregularly bedded sands, clays, and gravels, has an estimated thickness of 1,000 feet and rests unconformably on a basement of Paleozoic metamorphics, and in the east on Pre-Cambrian crystalline rocks and in part on Lower Cretaceous. The Eutaw formation, somewhat similar to the Tuscaloosa in lithologic character, is believed to be entirely marine, though much of the formation was doubtless laid down in very shallow water. It is 400-500 feet thick, rests conformably on the Tuscaloosa, and is overlain conformably in part by the Selma chalk, and in part by the Ripley formation. The Selma chalk consists mainly of more or less argillaceous and sandy limestones rendered chalky by their large content of foraminiferal remains. It is abundantly fossiliferous in certain portions, yielding large numbers of the *Exogyra* described in the latter portion of the paper. The Selma grades into the sandy member of the Tuscaloosa, and the clastic beds of the Ripley formation when followed along the strike. A thickness of 930 feet of the chalk formation has been measured in western Alabama. The Ripley formation, 250-350 feet in thickness, consists typically of calcareous and glauconitic sands, sandy clays, and impure limestones and marls of marine origin. It extends through parts of the Gulf states from southern Illinois to Georgia. A study of the faunas of the various formations is detailed, and correlations with other Cretaceous regions indicated by chart.

A description of the genus *Exogyra*, which includes three species with two varieties, constitutes the second portion of the paper.

---

R. C. M.

*The Jurassic Flora of Cape Lisburne, Alaska.* By F. H. KNOWLTON.

U.S. Geol. Surv., Prof. Paper 85, Part D, 1914. Pp. 25, pls. 4.

The Jurassic of the Cape Lisburne area is estimated to have a very great thickness, 15,000 feet, and contains from 40 to 50 coal beds which range in thickness from 1 or 2 feet to over 30 feet. Plant collections from this area show 17 species of well-defined Jurassic types. The close similarity or identity of a number of forms with species from eastern Siberia and Mongolia is noteworthy. The flora indicates a warm-temperate or subtropic climate and the geographic range, especially into the Arctic and Antarctic, is suggestive of the uniform mildness of the Jurassic earth-climate.

R. C. M.